

Table 1A

Cancer Incidence in Wayland, MA 1982-1992

Cancer Type	Total				Male				Female			
	Obs	Exp	SIR	95% CI	Obs	Exp	SIR	95% CI	Obs	Exp	SIR	95% CI
Bladder	22	24.6	90	56--136	19	18.2	104	63--163	3	6.4	NC	NC
Breast	97	94.2	103	83--126	1	0.6	NC	NC	96	93.6	103	83--125
Hodgkin's	5	4.8	104	33--243	2	2.6	NC	NC	3	2.2	NC	NC
Kidney	12	13.2	91	47--159	8	8.3	96	41--190	4	4.9	NC	NC
Leukemia	10	10.8	93	44--171	3	6.3	NC	NC	7	4.5	156	63--322
Liver	3	2.7	NC	NC	2	1.9	NC	NC	1	0.8	NC	NC
Lung	69	83.0	83	65--106	37	53.0	70 *	49--97	32	30.0	107	73--151
NHL	17	19.1	89	52--143	5	10.3	49	16--113	12	8.8	137	71--239
Pancreas	13	12.0	108	57--185	10	6.1	164	78--301	3	5.9	NC	NC
Stomach	7	12.2	57	23--118	5	7.8	57	23--118	2	4.5	NC	NC

Notes:

Expected number of cases presented are rounded to the nearest tenth. SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases Exp = Expected number of cases SIR = Standardized Incidence Ratio NC = Not Calculated

NHL = non-Hodgkin's Lymphoma Lung = Lung and Bronchus 95% CI = 95% Confidence Interval * Indicates statistical significance (P<.05)